

How dependent are the update of control and prediction during adaptation to visuomotor rotation?

2-F-109




James MATHEW¹, Frédéric DANION¹

¹ Institut de Neurosciences de la Timone, UMR 7289, CNRS-AMU, Campus Santé Timone, Marseille, France



INTRODUCTION

- **Prediction** turns motor commands into expected sensory consequences[1].
- **Control** turns desired consequences into motor commands.
- People can learn to predict the consequences of their actions before they can learn to control their actions [2].
- Update of prediction is faster than update of control [2].




- **GOAL** : to explore how the update of prediction might influence the update of control, and vice versa.
- **HOW** : by investigating the transfer of learning between two visuomotor tasks both requiring adaptation to a 90° rotation.

METHOD

- Two groups of 7 participants (mean age = 29.5, all right handed)

EXPERIMENTAL SETUP



PREDICTION TASK

- This task was used to probe the ability of participants to predict visual consequences arising from their hand actions
- Participants had to track with their eyes a self-moved cursor whose displacement was driven by a joystick [3].

CONTROL TASK

- This task was used to probe the ability to move a cursor with the hand along a specified 2D trajectory, which resulted from the combination of several sinusoids [4].
- Participants had to move the joystick so as to bring the cursor as close as possible from the moving target [5].

UPDATE in PREDICTION and CONTROL

- For both tasks, we introduced a 90° counter-clockwise visuomotor rotation altering the mapping between hand movements and their visual consequences on the screen (i.e. cursor motion).

EXPERIMENTAL DESIGN

Group 1

Predict 0° 10 trials (PRE)	Control 0° 10 trials (PRE)	Predict 90° 40 trials (learning)	Control 90° 40 trials (learning)	Control 0° 1 trial (POST)	Predict 0° 1 trial (POST)
----------------------------------	----------------------------------	--	--	---------------------------------	---------------------------------

Group 2

Control 0° 10 trials (PRE)	Predict 0° 10 trials (PRE)	Control 90° 40 trials (learning)	Predict 90° 40 trials (learning)	Predict 0° 1 trial (POST)	Control 0° 1 trial (POST)
----------------------------------	----------------------------------	--	--	---------------------------------	---------------------------------

Transfer is assessed through the group comparison




Fig. 1: Typical PREDICT trials from the same subject at various stages (only 2.5 sec are displayed out of 10s). Initial eye tracking is altered by the 90° rotation, but then improves across trials. Note also the after-effect.




Fig. 2: Typical CONTROL trials from the same subject at various stages of the experiment (only 2.5 sec are displayed out of 10s). Initial hand tracking is altered by the 90° rotation, but then improves across trials. Note also the after-effect.

RESULTS




Fig. 3: Positional error between eye and cursor in PREDICT and between cursor and target in CONTROL. Prior experience strongly reduced positional error in PREDICT, but not in CONTROL.




Fig. 5: Saccade rate during PREDICT and CONTROL tasks. Prior experience helped to somewhat reduce the saccade rate in PREDICT but not at all in CONTROL.




Fig. 4: Temporal lag between eye and cursor in PREDICT and between cursor and target in CONTROL. Prior experience tends to reduce the temporal lag in PREDICT, but not in CONTROL.






Fig. 6: Saccade amplitude during PREDICT and CONTROL tasks. Prior experience helped to strongly reduce the amplitude of saccade PREDICT, but not in CONTROL.

CONCLUSION

Preliminary results suggest the existence of an **asymmetrical transfer** such that prior experience in the CONTROL task benefited to performance in the PREDICT task, but not the other way around.



- ✓ A possible scheme to account for these results is that visuomotor adaptation in our CONTROL task requires both the update of a forward and an inverse model [6], whereas adaptation in our PREDICT task relies solely on the update of a forward model.



- ✓ At a more general level these results emphasize that our ability to predict movement consequences can be improved without necessarily improving our ability to control movement.

BIBLIOGRAPHY

- [1] Miall & Wolpert (1996) Neural Network 9: 1265-79
- [2] Flanagan et al (2003) Current Biology 13: 146-50
- [3] Landelle et al (2016) J Neurophysiology 116:1859-70
- [4] Soechting et al (2010) Plos One 5: e12574
- [5] Ogawa & Imamizu (2013) J Neuroscience 33:6412-22
- [6] Wolpert & Kawato (1998) Neural Network 11:1317-29

ACKNOWLEDGEMENT : This work was part of Innovative Training Network 'Perception and Action in Complex Environment' (PACE) that has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 642961 and also was supported by a French National Grant (REM ANR-13-APPR-0008).